

IJHSM

Indonesian Journal
on Health Science
and Medicine



UNIVERSITAS MUHAMMADIYAH SIDOARJO

Table Of Contents

Journal Cover	1
Author[s] Statement	3
Editorial Team	4
Article information	5
Check this article update (crossmark)	5
Check this article impact	5
Cite this article	5
Title page	6
Article Title	6
Author information	6
Abstract	6
Article content	8

Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

Indonesian Journal on Health Science and Medicine

Vol. 3 No. 1 (2026): July
DOI: 10.21070/ijhsm.v3i1.405

EDITORIAL TEAM

Editor in Chief

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia ([Google Scholar](#) | [Scopus ID: 57202239543](#))

Section Editor

Maria Istiqomah Marini, Department of Forensic Odontology, Faculty of Dentistry, Universitas Airlangga Surabaya, Indonesia ([Google Scholar](#) | [Scopus ID: 57214083489](#))

Heri Setiyo Bekti, Department of Medical Laboratory Technology, Poltekkes Kemenkes Denpasar, Indonesia ([Google Scholar](#) | [Scopus ID: 57194134610](#))

Akhmad Mubarok, Department of Medical Laboratory Technology, Universitas Al-Irsyad Al-Islamiyyah Cilacap, Indonesia ([Google Scholar](#))

Tiara Mayang Pratiwi Lio, Department of Medical Laboratory Technology, Universitas Mandala Waluya Kendari, Indonesia ([Google Scholar](#))

Syahrul Ardiansyah, Department of Medical Laboratory Technology, Faculty of Health Sciences, Universitas Muhammadiyah Sidoarjo, Indonesia ([Google Scholar](#) | [Scopus ID: 55390984300](#))

Miftahul Mushlih, Department of Medical Laboratory Technology, Faculty of Health Sciences, Universitas Muhammadiyah Sidoarjo, Indonesia ([Google Scholar](#) | [Scopus ID: 57215844507](#))


Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

Article information

Check this article update (crossmark)



Check this article impact (*)



Save this article to Mendeley



(*) Time for indexing process is various, depends on indexing database platform

Prevalence of Cryptosporidiosis in Preschool Children: Prevalensi Kriptosporidiosis pada Anak Usia Prasekolah

Baraa Abdulsalam Hraija Hraija, babdulsalam@uowasit.edu.iq (*)

Department of Microbiology, College of Medicine, University of Wasit, Wasit, Iraq

Dhamyaa Kareem Kadhim , thalsarraai@uowasit.edu.iq

Department of Anatomy and Biology, College of Medicine, University of Wasit, Wasit, Iraq

Ghasik Aqeel , galaqeeli@uowasit.edu.iq

Department of Microbiology, College of Medicine, University of Wasit, Wasit, Iraq

(*) Corresponding author

Abstract

General Background: Cryptosporidiosis is a globally distributed parasitic disease that frequently causes diarrheal illness in young children and represents a persistent public health concern. **Specific Background:** Cryptosporidium parvum is one of the most significant zoonotic protozoa responsible for gastrointestinal infections, and molecular identification methods are increasingly used to clarify epidemiological patterns and genetic diversity of circulating strains. **Knowledge Gap:** Despite numerous studies on intestinal parasites, molecular data regarding local isolates and subtype distribution of C. parvum in Iraqi children remain limited. **Aims:** This study aimed to detect C. parvum in diarrheic children using nested polymerase chain reaction targeting the GP60 gene and to analyze the genetic relationships of local isolates through sequencing and phylogenetic analysis. **Results:** Among 28 fecal samples examined, 12 (42.86%) were positive for C. parvum. Sequencing analysis of positive samples revealed a high genetic similarity (approximately 99%) with global reference strains in the NCBI database, with minimal nucleotide variation. Phylogenetic analysis further classified the detected isolates into two subtype groups, IIc and IIIId, with IIc representing the majority of cases. **Novelty:** This research provides the first molecular confirmation and phylogenetic characterization of these GP60 subtype groups among local Iraqi isolates deposited in the NCBI database. **Implications:** The findings contribute to the understanding of molecular epidemiology of cryptosporidiosis in Iraq and highlight the importance of expanded genotyping and surveillance studies to clarify transmission pathways and improve disease monitoring.

Keywords: Cryptosporidium Parvum, Molecular Epidemiology, Nested PCR, GP60 Gene, Pediatric Diarrhea

Key Findings Highlights

High proportion of pediatric stool samples contained detectable parasite DNA.

Genetic sequencing revealed strong similarity between local isolates and global strains.

Two allele groups dominated the detected variants within the sampled population.

Published date: 2026-03-05

Introduction

Vinayak et al. (2015) note that the diarrheal illnesses (particularly, the *Cryptosporidium* species, the first serious parasite organism that causes severe diarrhea) contributed to approximately 10.5% of all child deaths worldwide. Under the Eucoccidiorida Order of Apicomplexa Phylum, there is the *Cryptosporidium parvum* Family of opportunistic intracellular parasites (**Barta et al., 2012; Liu, 2017**). This parasite strikes a wide array of animals alongside people, triggering cryptosporidiosis (**Guérin and Striepen, 2020**). It ranks among the select parasites seeing rising incidence, with outbreaks increasingly routine nowadays (**Chalmers et al., 2019**).

The most common method of spreading the parasite by children below two years of age is through the consumption of infected water even though there are other methods through which the parasite is spread both directly and indirectly. *Cryptosporidium* can persist for long periods in the environment because its oocysts are protected by a thick outer shell, allowing them to withstand harsh conditions and many common disinfectants (**Robertson et al., 2020; Al-Ezzy and Kadhim, 2021**). Moreover, *Cryptosporidium* has the ability to proliferate in the small intestine microvilli, which disturbs ion homeostasis and leads to ionic loss in general (**Das et al., 2018; Mendes, 2020**). *Cryptosporidium* may also infect various sites throughout the gastrointestinal tract (**Peek et al., 2018**). Whereas, in several mild to moderate cases, cryptosporidiosis did not show any signs, severe cases may lead to vomiting, anorexia, fever, general malaise, abdominal cramping, and a great deal of watery diarrhea (**Peek et al., 2018**). The illness is common with a seroprevalence rate of 25-35% and infection rate of 1-2% across the world. The organism is capable of occurring in 1-4.5 percent of the sampled people (**Wanamaker and Grimm, 2004; Tulchinsky and Varavikova, 2014**). The organism was named blue beads due to the characteristic biopsy appearance of 1 to 5 mm-sized spherical and basophilic aggregates to appear out of the enterocytic apex in crypts or surface epithelium (**Jimenez et al., 2017; Schuetz, 2019**).

Traditional diagnostic approaches for detecting *Cryptosporidium* oocysts worldwide rely on light microscopy with Modified Ziehl-Neelsen staining, direct wet mounts, and ultrastructural techniques to visualize intracellular cysts (**Malik et al., 2013; Bones, 2017**). Nevertheless, due to different life cycle durations of *Cryptosporidium* strains and rather same appearance of this parasite, the conventional morphological and phenotypical systems cannot differentiate distinct species among humans and animals (**Wielinga et al., 2008**). Recent molecular diagnostic tools have refined *Cryptosporidium* species detection, strain subtyping, and genotyping, highlighting broad versus narrow host specificities in different strains (**Robinson and Chalmers, 2012**). Here, we deposited local positive *C. parvum* isolates into the NCBI database, classified their allelic profiles, and applied PCR-based methods to probe cryptosporidiosis at the molecular level.

Materials and Methods

Ethical approval

This study was allowed by the Scientific Committee of the College of Medicine, University of Wasit (Wasit, Iraq).

Study samples

This study enrolled 28 children with diarrhea from three government hospitals in Wasit Governorate, Iraq (Al-Kut Hospital for Gynecology, Obstetrics, and Pediatrics; Al-Karama Teaching Hospital; Al-Zahraa Teaching Hospital) between January and March 2022. Fresh stool samples were aseptically collected in disposable plastic containers from all participants, kept cool during transport, and processed for molecular analysis in the laboratory.

Molecular examination

DNA from stool samples was extracted following the manufacturer's protocol for the Stool DNA Extraction Kit (Bioneer, Korea). The concentration and purity of each of the DNA samples extracted were determined by the Nanodrop spectrophotometer. Nested PCR targeting the *GP60* gene, using two primer sets designed by **Maurya et al. (2013)**, confirmed via Primer3Plus and NCBI-GenBank, and synthesized by Bioneer (Korea), was employed to detect *C. parvum* (Table 1).

Primer	Sequence (5'-3')	Amplicon
First-step GP60 nested PCR forF	ATAGTCTCCGCTGTATTC	480bp
Cryptosporidium parvum R	GAGATATATCTTGGTGCG	

R CGAACCACATTACAAATGAAG

Table 1. **Table (1): Nested PCR primer sets targeting *C. parvum***

Nested PCR master mixes were prepared using the AccuPower[®] 2X PCR PreMix kit (Bioneer, Korea) in 20 µL reactions. The first round included 5 µL DNA template, 1 µL each of forward and reverse primers, and 13 µL PCR-grade water; the second round used 2.5 µL of the first-round product, 1 µL each primer, and 15.5 µL PCR-grade water. Thermal cycling (Thermocycler, Bioneer, Korea) consisted of initial denaturation at 95°C for 5 min, followed by 35 cycles of denaturation (95°C, 30 s), annealing (56°C, 30 s), and extension (72°C, 1 min), with a final extension at 72°C for 5 min. PCR products were resolved by electrophoresis on 2% agarose gels stained with ethidium bromide, visualized under UV transillumination,

and scored as positive for a 375 bp band

Phylogenetic analysis

The PCR products that were positive were transferred to Macogen Company (Korea) to undergo DNA sequencing in the modified Sanger method. The analysis of the results was then done. Local *C. parvum* strains received designated identifiers, were submitted to NCBI GenBank (with assigned accession numbers), and underwent NCBI-BLAST comparison with reference sequences for phylogenetic tree construction.

Results

A total of 28 fecal samples were collected and a nested PCR assay was performed on them; 12 of them (42.86) were positive in general (Figure 1). Ten genomic DNAs of positive samples were phylogenetically studied using the *GP60* gene. The name of the results of the sequencing of the local isolates of *Cryptosporidium parvum* was in the following way: Local cataloged variants—such as *C. parvum* Human/IQS-1, IQS-2, IQS-4, IQS-5, and IQS strains—displayed point mutations plus aligned matches across *GP60* segments (Figure 2)




Figure 1. **Figure (1): Agarose gel electrophoresis indicates the presence of the *GP60* gene of *C. parvum* in the feces of a human being when subjected to Nested PCR analysis.**

DNA Sequences		Translated Protein Sequences	
Species/Abbrv			
1. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.1	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
2. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.2	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
3. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.3	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
4. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.4	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
5. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.5	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
6. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.6	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
7. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.7	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
8. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.8	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
9. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.9	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
10. <i>Cryptosporidium parvum</i> isolate Human_IQS.No.10	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
11. AY738188.1 <i>Cryptosporidium parvum</i> isolate 7490	60	GTTCCTGACACACTCTCCAGTCCAGCAAGCACTTCCAGATGAAAGATGTCGGTACTTTCATTTTAAATGTTG	
12. AY873780.1 <i>Cryptosporidium parvum</i> 60-kDa glycoprote		GTTCCTGACACACTCTCCAGTCCAGCAAGCACTTCCAGATGAAAGATGTCGGTACTTTCATTTTAAATGTTG	
13. KX397568.1 <i>Cryptosporidium parvum</i> isolate BS98-Iic		GTTCCTGACACACTCTCCAGTCCAGCAAGCACTTCCAGATGAAAGATGTCGGTACTTTCATTTTAAATGTTG	
14. DQ192508.1 <i>Cryptosporidium parvum</i> isolate 9837	60	GTTCCTGACACACTCTCCAGTCCAGCAAGCACTTCCAGATGAAAGATGTCGGTACTTTCATTTTAAATGTTG	
15. AY166805.1 <i>Cryptosporidium parvum</i> isolate 4736	60	GTTCCTGACACACTCTCCAGTCCAGCAAGCACTTCCAGATGAAAGATGTCGGTACTTTCATTTTAAATGTTG	
16. FJ839876.1 <i>Cryptosporidium parvum</i> isolate MQ153	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
17. KX397563.1 <i>Cryptosporidium parvum</i> isolate BS7-IIId	60	CTGCTGAAAGGAGCCCGAAACTCCCGAATCTTACTCCAAAGGAAAGATGTTGGTACTTTCATTTTAAATGTTG	
18. AY382675.1 <i>Cryptosporidium parvum</i> isolate A62	60kDa	GTTCCTGACACACTCTCCAGTCCAGCAAGCACTTCCAGATGAAAGATGTCGGTACTTTCATTTTAAATGTTG	

Figure 2. Lane M: DNA marker ladder (2000–100 bp); Lanes 1–12: Positive amplicons at 375 bp; NTC: No-template control

The IQS-*C. parvum* isolates 2, 5, and 9 showed significant sequence identity with the Egyptian *C. parvum* isolate (KX397563.1), as determined by NCBI-BLAST homology analysis. All local *C. parvum* subtypes subsequently clustered into the Iic (7/10 isolates) and IIId (3/10 isolates) allele groups (Figure 3).




Figure 3. **Figure (3): BLAST sequence identity (%) distribution across local *C. parvum* isolates compared to reference strains**

The local isolates and global isolates were compared and it was observed that the total genetic mutation of the local *C. parvum* IQS-isolates was 0-0.9% and that it had a high similarity (99% similarity) to the NCBI-BLAST *C. parvum* of the gene GP60 (Figure 4).




Figure 4. **Figure (4): The comparison of the NCBI-GenBank isolates using the GP60 gene with the incomplete sequences of local *C. parvum* IQS isolates using phylogenetic tree.**

Discussion

A close relationship between *Cryptosporidium* and both acute and chronic diarrhea in children has been demonstrated in a wide range of countries. In Iraq, investigations have largely depended on classic approaches like In Iraq, conventional microscopy via modified Ziehl-Neelsen (Alali et al., 2021) has dominated, offering scant molecular evidence. PCR-based *C. parvum* rates among diarrheal youth: Baghdad 11% (Hussein et al., 2015), Al-Muthanna 18% (Mallah & Jomah, 2015), Al-Diwaniyah 24% (Ahmed et al., 2016), Erbil 12% (Azeez & Alsakee, 2017), Al-Najaf 12.8% (Tairsh et al., 2017), Thi-Qar 10.42% (Salim & Al-Aboody, 2019). Comparable figures abroad included 10% (Netherlands), 3.77% (Ethiopia), 10.42% (Brazil), and 7.14% (Turkey) (Wielinga et al., 2008; Adamu et al., 2010; Taghipour et al., 2011; Rolando et al., 2012; Yilmazer et al., 2017).

Discrepancies between our findings and other local/international studies may stem from differences in targeted genes, seasonal variations in cryptosporidiosis incidence, sampling biases (sample size, selection methods, patient age), environmental parasite sources, and PCR conditions. Global reports on *Cryptosporidium* prevalence in children vary widely, with *C. hominis* predominating in South Africa (Leav et al., 2002), Thailand (Tiangtip and Jongwutiwes, 2002), Malawi (Peng et al., 2003), Brazil (Bushen et al., 2007), Kenya (Mbae, 2008), Peru (Cama et al., 2008), and South India (Ajjampur et al., 2010), whereas *C. parvum* was the main species in Kuwait (Sulaiman et al., 2005), Ethiopia (Adamu et al., 2010), Iran (Taghipour et al., 2011), and Turkey (Yilmazer et al., 2017).

Genotyping and subtyping efforts for *Cryptosporidium* routinely examine markers like 18S rRNA, the 70-kDa heat shock protein (hsp70), oocyst wall protein (OWP), actin, β -tubulin, TRAP, ITS1, and DHFR (Cunha et al., 2019). *GP60* remains a premier target for *C. parvum* subtype discrimination (Khan et al., 2018; Yanta et al., 2021; Uran-Velasquez et al., 2022).

Strains with identical *GP60* genotypes can vary substantially at additional genetic loci, sometimes exceeding differences seen between distinct *GP60* alleles (Abal-Fabeiro et al., 2013). The Ic allele, initially identified in *C. hominis*, has been found exclusively in human *C. parvum* bovine genotype isolates (Alves et al., 2003). The Iic subtype is enriched in short 9-serine repeats and is rare in animals, possibly due to its wide geographic distribution and partial overlap in human-animal sequence origins (Widmer et al., 2009).

The IId subtype is uncommon among *C. parvum* strains but linked to zoonotic cases in European countries including Italy, Hungary, Portugal, and Serbia (Xiao and Fayer, 2008), comprising about half of pediatric infections in Kuwait (Sulaiman et al., 2005). Variations in short tandem repeats and host immune responses may impose differing selective pressures on *GP60* across species, favoring short-repeat alleles in humans (Widmer et al., 2009).

Conclusion

In conclusion, this study revealed a notably high prevalence of *C. parvum* in Iraqi children with diarrhea and provided the first confirmation in Iraq of allelic group distributions (Iic and IId) among local isolates. The transmission sources and pathways for *C. parvum* in Iraq have yet to be clarified. Additional studies focusing on serotyping and genotyping *Cryptosporidium* species among patients with diarrhea are crucial to address these informational voids.

Authors' Contributions

DKK gathered fecal specimens from diarrheic pediatric patients and carried out DNA isolation. BAH and GA handled the nested PCR experiments and phylogenetic evaluations. All authors contributed to genotyping the positive isolates and drafting the manuscript.

Competing Interests

There is no competing to be interested, and no funds have received to complete this work.

References

- [1] J. L. Abal-Fabeiro, X. Maside, X. Bello, J. Llovo, and C. Bartolome, "Multilocus Patterns of Genetic Variation Across *Cryptosporidium* Species Suggest Balancing Selection at the GP60 Locus," *Molecular Ecology*, vol. 22, no. 18, pp. 4723-4732, 2013.
- [2] H. Adamu, B. Petros, A. Hailu, and F. Petry, "Molecular Characterization of *Cryptosporidium* Isolates From Humans in Ethiopia," *Acta Tropica*, vol. 115, no. 1-2, pp. 77-83, 2010.
- [3] H. S. Ahmed, A. H. Abd, and N. Q. Mohammed, "Detection of *Cryptosporidium parvum* From Feces Samples of Human and Camels Using Direct PCR Technique," *Al-Qadisiyah Journal of Veterinary Medicine Sciences*, vol. 15, no. 2, pp. 59-62, 2016.
- [4] S. S. R. Ajjampur et al., "Multisite Study of Cryptosporidiosis in Children With Diarrhea in India," *Journal of Clinical Microbiology*, vol. 48, no. 6, pp. 2075-2081, 2010.
- [5] F. Alali, I. Abbas, M. Jawad, and N. Hijjawi, "Cryptosporidium Infection in Humans and Animals From Iraq A Review," *Acta Tropica*, vol. 220, 2021.
- [6] A. I. A. Al-Ezzy and A. T. Kadhim, "Evaluation for Sociodemographic Risk Factors Associated With *Cryptosporidium parvum* Infection in Children," *Diyala Journal for Veterinary Sciences*, vol. 1, no. 2, pp. 100-113, 2021.
- [7] M. Alves et al., "Subgenotype Analysis of *Cryptosporidium* Isolates From Humans and Animals," *Journal of*

Clinical Microbiology, vol. 41, no. 6, pp. 2744-2747, 2003.

8. [8] S. S. Azeez and H. M. Alsakee, "Cryptosporidium spp. and Rotavirus Gastroenteritis in Children," *Medical Journal of Indonesia*, vol. 26, no. 3, pp. 190-197, 2017.
9. [9] J. R. Barta et al., "Phylogenetic Position of Adeleorinid Coccidia Inferred Using 18S rDNA Sequences," *Journal of Eukaryotic Microbiology*, vol. 59, no. 2, pp. 171-180, 2012.
10. [10] A. J. Bones, *Developing in Vitro Culturing Techniques of Cryptosporidium parvum*, University of Kent, 2017.
11. [11] O. Y. Bushen et al., "Heavy Cryptosporidial Infections in Children in Brazil," *Transactions of the Royal Society of Tropical Medicine and Hygiene*, vol. 101, no. 4, pp. 378-384, 2007.
12. [12] V. A. Cama et al., "Cryptosporidium Species and Clinical Manifestations in Children," *Emerging Infectious Diseases*, vol. 14, no. 10, p. 1567, 2008.
13. [13] R. M. Chalmers, G. Robinson, K. Elwin, and R. Elson, "Analysis of Cryptosporidium Subtypes Linked to Human Outbreaks," *Parasites & Vectors*, vol. 12, no. 1, 2019.
14. [14] F. S. Cunha, R. H. S. Peralta, and J. M. Peralta, "Detection and Molecular Characterization of Cryptosporidium," *Revista do Instituto de Medicina Tropical de Sao Paulo*, vol. 61, 2019.
15. [15] S. Das, R. Jayaratne, and K. E. Barrett, "Role of Ion Transporters in Infectious Diarrhea," *Cellular and Molecular Gastroenterology and Hepatology*, vol. 6, no. 1, pp. 33-45, 2018.
16. [16] A. Guérin and B. Striepen, "Biology of the Intestinal Parasite Cryptosporidium," *Cell Host and Microbe*, vol. 28, no. 4, pp. 509-515, 2020.
17. [17] R. A. Hussein et al., "Evaluation of Multiplex Real-Time PCR for Intestinal Parasites," *International Journal*, vol. 3, no. 9, pp. 782-788, 2015.
18. [18] M. Jimenez, B. Miller, and H. L. Bridle, "Separation of Microparticles for Waterborne Pathogens," *Chemical Engineering Science*, vol. 157, pp. 247-254, 2017.
19. [19] A. Khan, J. S. Shaik, and M. E. Grigg, "Genomics and Molecular Epidemiology of Cryptosporidium Species," *Acta Tropica*, vol. 184, pp. 1-14, 2018.
20. [20] B. A. Leav et al., "Sequence Diversity at the Cpgp40/15 Locus Among Cryptosporidium Isolates," *Infection and Immunity*, vol. 70, no. 7, pp. 3881-3890, 2002.
21. [21] D. Liu, "Cryptosporidium," in *Laboratory Models for Foodborne Infections*, CRC Press, 2017, pp. 589-597.
22. [22] A. Malik, J. Kulaar, R. Shukla, and V. Dutta, "Triple Protozoal Enteropathy of the Small Intestine," *Indian Journal of Sexually Transmitted Diseases and AIDS*, vol. 34, no. 2, p. 119, 2013.
23. [23] M. O. Mallah and N. R. Jomah, "Epidemiological and Molecular Study of Cryptosporidium parvum," *Donnish Journal of Microbiology and Biotechnology Research*, vol. 2, no. 1, pp. 1-7, 2015.
24. [24] P. S. Maurya et al., "Genotyping of Cryptosporidium Species in Bovine Calves," *Indian Journal of Animal Sciences*, vol. 83, pp. 1018-1023, 2013.
25. [25] C. Mbae, "Cryptosporidiosis Prevalence and Genotype Analysis," *International Journal of Infectious Diseases*, vol. 12, p. e72, 2008.
26. [26] B. D. P. Mendes, *Etiology of Neonatal Diarrhea in Calves in Germany*, Universidade de Lisboa, 2020.
27. [27] S. F. Peek et al., "Infectious Diseases of the Gastrointestinal Tract," in *Rebhun's Diseases of Dairy Cattle*, 2018.
28. [28] M. M. Peng et al., "Molecular Epidemiology of Cryptosporidiosis in Children in Malawi," *Journal of Eukaryotic Microbiology*, vol. 50, pp. 557-559, 2003.
29. [29] L. J. Robertson et al., "Cryptosporidium Infections in Africa," *Frontiers in Veterinary Science*, vol. 7, 2020.
30. [30] G. Robinson and R. M. Chalmers, "Assessment of Genetic Markers for Typing Cryptosporidium," *Experimental Parasitology*, vol. 132, no. 2, pp. 200-215, 2012.
31. [31] R. F. R. Rolando et al., "Detection and Differentiation of Cryptosporidium by Real-Time PCR," *Memorias do Instituto Oswaldo Cruz*, vol. 107, pp. 476-479, 2012.
32. [32] A. R. Salim and B. A. Al-Aboody, "Molecular Detection of Cryptosporidium parvum," *Iraqi Journal of Biotechnology*, vol. 18, no. 2, 2019.
33. [33] A. N. Schuetz, "Infectious Disorders of the Duodenum and Small Bowel," in *Surgical Pathology of Non-Neoplastic Gastrointestinal Diseases*, 2019.
34. [34] I. M. Sulaiman et al., "Endemicity of Cryptosporidiosis in Children in Kuwait," *Journal of Clinical Microbiology*, vol. 43, no. 6, pp. 2805-2809, 2005.
35. [35] N. Taghipour et al., "Molecular Epidemiology of Cryptosporidiosis in Iranian Children," *Iranian Journal of Parasitology*, vol. 6, no. 4, p. 41, 2011.
36. [36] H. R. Tairsh et al., "Identification of Cryptosporidium spp. Infections by PCR Technique," *European Journal of Pharmaceutical and Medical Research*, vol. 4, no. 2, pp. 208-215, 2017.
37. [37] R. Tiangtip and S. Jongwutiwes, "Molecular Analysis of Cryptosporidium Species in HIV Patients," *Tropical Medicine and International Health*, vol. 7, no. 4, pp. 357-364, 2002.
38. [38] T. H. Tulchinsky and E. A. Varavikova, *The New Public Health*, Academic Press, 2014.
39. [39] J. Uran-Velasquez et al., "Multilocus Sequence Typing of Cryptosporidium," *PLOS ONE*, vol. 17, no. 7, 2022.
40. [40] S. Vinayak et al., "Genetic Modification of Cryptosporidium parvum," *Nature*, vol. 523, pp. 477-480, 2015.
41. [41] R. Wanamaker and I. Grimm, *Encyclopedia of Gastroenterology*, 2004.
42. [42] G. Widmer, "Meta-Analysis of a Surface Glycoprotein of Cryptosporidium," *Epidemiology and Infection*, vol. 137, no. 12, pp. 1800-1808, 2009.
43. [43] P. R. Wielinga et al., "Molecular Epidemiology of Cryptosporidium in Humans and Cattle," *International Journal for Parasitology*, vol. 38, no. 7, pp. 809-817, 2008.
44. [44] L. Xiao and R. Fayer, "Molecular Characterisation of Cryptosporidium and Giardia," *International Journal for Parasitology*, vol. 38, no. 11, pp. 1239-1255, 2008.
45. [45] C. A. Yanta et al., "CryptoGenotyper Bioinformatics Tool for Cryptosporidium Identification," *Food and Waterborne Parasitology*, vol. 23, 2021.
46. [46] N. Yilmazer et al., "Cryptosporidiosis in Humans With Reference to the First Case of Cryptosporidium hominis

Indonesian Journal on Health Science and Medicine

Vol. 3 No. 1 (2026): July
DOI: 10.21070/ijhsm.v3i1.405

Infection in Turkey," Medical Bulletin of Haseki, vol. 55, pp. 194-198, 2017.